

CHILD/ADOLESCENT MEASURES (0 through 18 years)	GUIDELINE	HEDIS COMPLIANT CPT/ICD9 CODES DOCUMENTATION TIPS
Well Care		
Access To PCP	Ensure a preventive/ambulatory care PCP visit at least annually.	Visit: ICD-9: V20.2, V70.0, V70.3, V70.5, V70.6, V70.8, V70.9 CPT: 99201-99205, 99211-99215, 99241-99245; 99381-99385, 99391-99395, 99401-99404, 99411, 99412, 99420, 99429
Adolescent Preventive Care	<p>Document risk assessment screening or counseling for adolescents 12-17yrs old annually:</p> <ol style="list-style-type: none"> 1. Sexual activity 2. Depression 3. Tobacco Use 4. Substance Use <p><i>These 4 topics are in addition to the 3 topics listed under Well Child Visits (health and development history, mental and physical), physical exam, health education/anticipatory guidance) for a total of 7 required screening/counseling indicators.</i></p>	<p>This is a 100% medical record review measure.</p> <p>Related sexual activity codes: ICD9- V65.44, V65.44, V25.0-V25.04, V25.09 Depression Screening: V79.0 Tobacco cessation counseling and classes: CPT: 99406, 99407; HCPCS: S9453</p> <p>If using acronyms for assessment/counseling tools, such as HEADS or TIPS, make sure all issues addressed are clearly identified/spelled out.</p> <p>Likewise, stating "Toxic Habits" is insufficient without identifying specific topics covered within that heading.</p> <p>A self administered questionnaire is a good way to cover all required topics, but the answers themselves must be in the chart. It is not sufficient to state "GAPS/ mini-GAPS completed"</p> <p>A sticker or stamp checklist for all topics is a good documentation tool.</p>

Chlamydia Screening	Screen sexually active females 16-24 yrs old annually.	Chlamydia Screen: CPT: 87110,87270,87320,87490,87491,87492,87810 Do not use CPT code 87800 for Chlamydia screening; it is not specific to Chlamydia and is not HEDIS-compliant.
CHILD/ADOLESCENT MEASURES (0 through 18 years)	GUIDELINE	HEDIS COMPLIANT CPT/ICD9 CODES DOCUMENTATION TIPS
Childhood Immunizations	Complete immunizations on or before the child's 2nd birthday: 4 DTaP/DTP 3 IPV 3 Hib 3 Hep B 4 PCV 1 MMR 1 VZV 3 Rotavirus 1 Hep A 2 Influenza	NYC PCPs must report all immunizations electronically through the Citywide Immunization Registry. www.nyc.gov/html/doh/html/cir/a01.html DtaP CPT: 90698,90700,90721, 90723 IPV CPT: 90698, 90713, 90723 Hib CPT: 90645-90648,90698,90721,90748 Hep B CPT: 90723, 90740, 90744, 90747, 90748 PCV CPT: 90669, 90670 MMR CPT: 90707, 90710 VZV CPT: 90710, 90716 RV CPT: 2 dose: 90681, 3 dose: 90680 Hep A CPT: 90633 Flu CPT: 90655, 90657, 90661, 90662
Adolescent Immunizations	Complete immunizations on or before the child's 13th birthday: 1 MCV4 or MPSV4. 1 Tdap or TD	MCV4 CPT: 90734; MPSV4 CPT: 90733 Tdap CPT: 90715; Td CPT: 90718
Dental Visit	Refer children 2-18 yrs old to the dentist annually.	Healthplex is Affinity's dental benefit manager. Call 1-800-468-9868 or go to www.Healthplex.com for a list of dental providers. For a list of preferred dentists in your neighborhood, contact Affinity's QM Department at 1-718-794-6034.
Human Papillomavirus Vaccine for Female Adolescents (HPV)	Ensure that female adolescents receive three doses of HPV vaccine by their 13 th birthday	CPT: 90649, 90650
Lead Testing	Test by 12 months and again by 24 months of age.	Lead Screen: CPT: 83655
Well Child Visits (WCV):		NB: The rendering of a preventive service at any sick or follow-up visit warrants the use of the preventive CPT E&M codes listed below.
0-15 Months	Ensure 6 or more well child visits by 15 months of age	Visit: ICD-9 Diag: V20.2, V20.3, V70.0, V70.3, V70.5, V70.6, V70.8, V70.9 with E&M codes CPT: 99381,99382,99391,99392, 99461

3- 6 Years Old	Ensure 1 well child visit annually	Visit: ICD-9 Diag: V20.2, V70.0, V70.3, V70.5, V70.6, V70.8, V70.9 with E&M codes CPT: 99382,99383,99392,99393
12-21 Years Old	Ensure 1 well child visit annually	Visit: ICD-9 Diag: V20.2, V70.0, V70.3, V70.5, V70.6, V70.8, V70.9 with E&M codes CPT: 99383-99385,99393-99395 Do not use ICD-9 Diagnosis V21.2 for adolescent well visits; it does not connote well care and is not HEDIS-compliant.
CHILD/ADOLESCENT MEASURES (0 through 18 years)	GUIDELINE	HEDIS COMPLIANT CPT/ICD9 CODES DOCUMENTATION TIPS
Weight Assessment and Counseling for Nutrition and Physical Activity	Document counseling/assessment for children 3-17 yrs annually: 1. BMI percentile * 2. Nutrition 3. Physical Activity * Height and weight must be documented in medical record	BMI Percentile: ICD-9 Proc: V85.5 Nutrition Counseling: ICD-9: V65.3; CPT: 97802-97804 Physical Activity Counseling: ICD-9: V65.41; HCPCS: S9451
Chronic Care		
Follow-up Care for Children with ADHD Medications	Ensure 3 follow up visits in 10 months for children 6-12 yrs old with newly prescribed ADHD medication: 1. visit within 30 days of initial dispensing of medication 2. additional visits between 31 days and 10 months after initial dispensing of medication. One visit can be a telephone visit.	Telephone Visit E&M Codes: CPT: 98966-98968, 99441-99443
HIV/AIDS Comprehensive Care	For HIV + patients between 2 and 18 years old: 1. Conduct visits at least twice a year, with 1 visit occurring <u>between 1/1 and 6/30</u> , and 1 visit occurring <u>between 7/1 and 12/31</u> (one in the first 6 months; one in the second 6 months) 2. Test viral load at least twice a year, with 1 test occurring <u>between 1/1 and 6/30</u> , and 1 test occurring <u>between 7/1 and 12/31</u> (one in the first 6 months; one in the second 6 months)	Visit: ICD9: V20.2,V70.0,V70.3,V70.5,V70.6, V70.8, V70.9 Viral Load Test: CPT: 87534-87536

Use of Appropriate Medications for Asthma: 3 Controllers	Prescribe a controller for children 5-18 yrs old at least 3 times annually. Preferred therapies: Cromolyn sodium inhaled corticosteroids, leukotriene modifiers, methylxanthines, nedocromil. Add on therapy: long-acting inhaled beta-2 agonists	ICD9: 493.0, 493.1, 493.8, 493.9 Inhaled short acting beta agonists, systemic corticosteroids, and oral and nasal preparations are not HEDIS-compliant medications.
Medication Management for People with Asthma	Follow patients 5-64 yrs prescribed asthma controller medications to ensure they are filling their prescriptions during the treatment period, i.e., the period of time beginning on the IPSD through the last day of the measurement year. Two rates are reported: 1. The percentage of members who remain on the controller medications at least 50% of their treatment period. 2. The percentage of members who remain on the controller medications at least 75% of their treatment period	ICD9: 493.0, 493.1, 493.8, 493.9 Inhaled short acting beta agonists, systemic corticosteroids, and oral and nasal preparations are not compliant for controller medication prescribing
Asthma Medication Ratio	Patients 5-64 yrs diagnosed with persistent asthma should be treated <u>preferably with controller medications</u> . The proportion of controller medications to all medications prescribed during the measurement year to treat asthma should be greater than 50%.	CD9: 493.0, 493.1, 493.8, 493.9 Inhaled short acting beta agonists, systemic corticosteroids, and oral and nasal preparations are not compliant for controller medication prescribing.
Acute Care		
Appropriate Treatment for Children with Pharyngitis	Perform a Rapid Strep test/throat culture prior to prescribing antibiotics for children/adolescents 2-18 yrs old with pharyngitis. <i>1. Diagnosis must be of <u>only pharyngitis</u> to be counted.</i> <i>2. Antibiotics prescribed <u>on or up to 3 days after</u> the diagnosis of pharyngitis are counted.</i>	ICD9: Acute Pharyngitis: 462 Acute Tonsillitis 463 Streptococcol sore throat: 034.0 Rapid StrepTest: CPT: 87880 Culture: CPT: 87070,87071,87081,87430,87650-87652 Rapid Strep Tests are billable; Remember to submit claims
Appropriate Treatment for Children with Upper Respiratory Infection (URI)	Do not <u>prescribe antibiotics</u> for children 3 months to 18 years old diagnosed with <i>only</i> URI	URI: ICD9: 460,465 Prescribe antibiotics only if you document an additional and/or different diagnosis.